

Evidence of a false thumb in a fossil carnivore clarifies the evolution of pandas

Manuel J. Salesa*[†], Mauricio Antón*, Stéphane Peigné[‡], and Jorge Morales*

*Departamento de Paleobiología, Museo Nacional de Ciencias Naturales, Consejo Superior de Investigaciones Científicas, José Gutiérrez Abascal 2, 28006 Madrid, Spain; and [‡]Laboratoire de Géobiologie, Biochronologie et Paléontologie Humaine, Centre National de la Recherche Scientifique Unité Mixte de Recherche 6046, Université de Poitiers, 40 Avenue du Recteur Pineau, 86022 Poitiers, France

Edited by F. Clark Howell, University of California, Berkeley, CA, and approved October 7, 2005 (received for review June 12, 2005)

The “false thumb” of pandas is a carpal bone, the radial sesamoid, which has been enlarged and functions as an opposable thumb. If the giant panda (*Ailuropoda melanoleuca*) and the red panda (*Ailurus fulgens*) are not closely related, their sharing of this adaptation implies a remarkable convergence. The discovery of previously unknown postcranial remains of a Miocene red panda relative, *Simocyon batalleri*, from the Spanish site of Batallones-1 (Madrid), now shows that this animal had a false thumb. The radial sesamoid of *S. batalleri* shows similarities with that of the red panda, which supports a sister-group relationship and indicates independent evolution in both pandas. The fossils from Batallones-1 reveal *S. batalleri* as a puma-sized, semiarboreal carnivore with a moderately hypercarnivore diet. These data suggest that the false thumbs of *S. batalleri* and *Ailurus fulgens* were probably inherited from a primitive member of the red panda family (Ailuridae), which lacked the red panda’s specializations for herbivory but shared its arboreal adaptations. Thus, it seems that, whereas the false thumb of the giant panda probably evolved for manipulating bamboo, the false thumbs of the red panda and of *S. batalleri* more likely evolved as an aid for arboreal locomotion, with the red panda secondarily developing its ability for item manipulation and thus producing one of the most dramatic cases of convergence among vertebrates.

convergence | *Simocyon* | Miocene | Batallones-1 | Ailuridae

The fossil site of Batallones-1 (Madrid) is a Miocene carnivore trap, which is greatly improving our knowledge about the Late Miocene European carnivores (1, 2). With 98% of the remains corresponding to members of Carnivora and an unbiased representation of skeletal parts, the site is yielding unprecedented information about the anatomy of species that previously were poorly known. One example is *Simocyon batalleri*, traditionally known from cranial fragments and dentition, which is now the best-represented member of the genus, thanks to the Batallones-1 material. A study of the cranial fossils from Batallones-1 has provided further evidence of the systematic position of *Simocyon* as a member of the Ailuridae, has shown a trend in the evolution of this genus toward increased hypercarnivory and bone crushing, and has typified *S. batalleri* as a hunter of small prey and occasional scavenger (3). Now, the discovery that this animal had a “false thumb” throws new light on the evolution of this feature in both the giant panda (*Ailuropoda melanoleuca*) and the red panda (*Ailurus fulgens*) lineages.

S. batalleri is the least abundant carnivore in the Batallones-1 association, and the fossils recovered correspond to two individuals. The fossils include crania and mandibles, cervical, thoracic, and lumbar vertebrae, nearly complete forelimbs, and partial hindlimbs (Fig. 1). The cervical vertebrae are unspecialized and similar to those of *Ailurus fulgens*, but the lumbar vertebrae are elongated, with pointed neural processes and elongated transverse processes, betraying a powerful musculature for flexing the back and the femur. These features are adaptations for a bounding gallop, a gait common in small musteloid carnivores, such as weasels (4), but energetically costly

and thus uncommon in larger, terrestrial carnivores. *Simocyon*, about the size of a puma, is the largest known arctoid carnivore with such lumbar morphology, likely reflecting a scansorial habit, meaning that the animal would forage on the ground but would readily climb when necessary (5). Many features of the appendicular skeleton indicate climbing abilities and the lack of cursorial adaptations. The scapula has a large process for the teres major muscle, shared only with the arboreal kinkajou (*Potos flavus*) among the musteloidea and superficially similar to that of bears, which combines a large teres major process with the development of a postscapular fossa, absent in *Simocyon* (5). The morphology of the long bones of the forelimb indicates the presence of strong flexor muscles in addition to a great ability for lateral rotation and a crouched, semiplantigrade posture (5).

The radial sesamoid of *Simocyon* (Fig. 2) is the only known instance of a well developed false thumb in any carnivore, living or fossil, other than the pandas. It is elongated, with an articulation facet for the posterior tubercle of the scapholunar; this facet is posterolaterally placed in the giant panda, whereas, in *Simocyon* and the red panda, it faces laterally. The medial margin of the bone in the three species shows three small facets for the attachment of several ligaments and the abductor pollicis brevis and opponens pollicis muscles, whereas, in the lateral surface, there is a large attachment area for the abductor pollicis longus muscle. Although both pandas have hypertrophied radial sesamoids, their morphologies are different: in the giant panda, it is relatively much larger and flattened, with a rough and massive tip, whereas, in the red panda and *Simocyon*, it is relatively smaller and not flattened, with a concave tip. An additional difference in the anatomy of the manus is that, in the red panda (and all procyonids), there is a flexor brevis digitorum manus muscle attaching on the fifth metacarpal that flexes the hand and reinforces the grasping action of the radial sesamoid, whereas the giant panda lacks this muscle (6, 7). As in modern pandas, the false thumb of *Simocyon* would be associated with a more or less differentiated section of the palmar pad, and the pulling action of the abductor–adductor musculature on the bone would fold the pad, allowing it to close around objects. This mechanism provides gripping abilities to a hand otherwise constrained by the lack of opposability of the true thumb or pollex, which is aligned with the other digits. Pandas employ this gripping action for grasping and manipulating bamboo stems and leaves (6–10), but this gripping action would also improve the grip of the forepaws of *Simocyon* during arboreal locomotion, especially on thin branches. Independently of the evolutionary causes of the origin of the false thumb, such arboreal adaptations would be important for a medium-sized carnivore, with small canines and without cursorial abilities like *Simocyon*, which coexisted with larger, dangerous competitors like the saber-toothed cats, *Machairodus* and *Paramachairodus*, and the bear-dog *Amphi-*

Conflict of interest statement: No conflicts declared.

This paper was submitted directly (Track II) to the PNAS office.

[†]To whom correspondence should be addressed. E-mail: mcns188@mncn.csic.es.

© 2005 by The National Academy of Sciences of the USA




Fig. 1. Skeletal anatomy of *S. batalleri*. (A) Skeletal reconstruction of *S. batalleri*. The pelvis, femora, tibiae, fibulae, sacrum, and caudal vertebrae are not known and have been reconstructed on the basis of related taxa. (B) Articulated right carpus and metacarpus in palmar view, showing the position of the radial sesamoid (rs) in *Ailuropoda melanoleuca* (Left; 1) and *S. batalleri* (Right; 2) (not at scale). Art by M. Antón; manus of giant panda modified from ref. 6.

cyon. Both *Machairodus* and *Amphicyon* were much larger than *Simocyon*, but their large size limited their climbing abilities, so it would be relatively simple for *Simocyon* to escape from them

by using short bursts of speed that would take it to the nearest tree and then climbing to safety. Confrontation with *Paramachairodus* would pose a different problem because this puma-sized saber-toothed cat was both dangerously armed and a capable climber. An ability to move along thin branches would have allowed *Simocyon* both to access carcasses cached by *Paramachairodus* on the trees and to retreat beyond the reach of even this agile cat.

This evidence has implications for the relationship between the giant and red pandas and for the evolution of their false thumbs. Although most specialists now agree about the ursid affinities of the giant panda (11–14), the position of the red panda has remained controversial. Molecular and morphological evidence has been interpreted by some to support a sister-group relationship with the Musteloidea (Procyonidae + Mustelidae) as the family Ailuridae (13–17), whereas others include it in the Procyonidae, as the subfamily Ailurinae (18, 19). Still others have proposed a close relationship between the red panda and the ursids, or ursids + pinnipeds (20–22).

The similarities in the radial sesamoids of *Simocyon* and *Ailurus* reinforce the notion of a sister-group relationship between the two genera, to the exclusion of procyonids and other musteloids to which the ailurids have been previously associated (Fig. 3). This fact agrees with views that postulate an independent family Ailuridae as a basal sister-group of Musteloidea (13–17). In all known carnivores, other than the pandas and *Simocyon*, the radial sesamoid retains its primitive state as a small, lens-shaped bone that develops on the tendon of the abductor pollicis longus muscle where it glides over the scapholunar. In turn, the false thumb of the giant panda has been seen as an exaggeration of a trend toward hypertrophy of the radial sesamoid already present in ursids, which also have the same distinctive pattern of muscle insertions (6). In summary, the retention of the primitive pattern in all other arctoids and the morphological differences between ailurids and the giant panda




Fig. 2. Comparisons of left radial sesamoids of the three species of carnivores with false thumbs, represented at the same size. (Left) Internal face. (Right) External face. Am, *Ailuropoda melanoleuca*; Sb, *Simocyon batalleri*; Af, *Ailurus fulgens*; 1, abductor pollicis brevis muscle; 2, abductor pollicis longus muscle; 3, opponens pollicis muscle; a, articulation facet with scapholunar.

We thank J. F. Pastor, from the Museo Anatómico (Valladolid, Spain), for access to osteological specimens. We also thank the Comunidad Autónoma de Madrid (Dirección General de Patrimonio Histórico) for the continuous funding support and research permissions. This study is part of Research Projects BTE2002-00410 (to J.M.) (Dirección General

de Investigación; Ministerio de Ciencia y Tecnología) and EX2003-0243 (to M.J.S.) (Secretaría de Estado de Educación y Universidades and European Social Funding). National Geographic Society Grant 6964-01 (to J.M.) and the Fondation Singer-Polignac (Paris) provided additional support.

1. Pozo, M., Calvo, J. P., Silva, P., Morales, J., Peláez-Campomanes, P. & Nieto, M. (2004) *Geogaceta* **35**, 143–146.
2. Morales, J., Alcalá, L., Álvarez-Sierra, M. A., Antón, M., Azanza, B., Calvo, J. P., Carrasco, P., Fraile, S., García-Paredes, I., Gómez, E., *et al.* (2004) *Geogaceta* **35**, 139–142.
3. Peigné, S., Salesa, M. J., Antón, M. & Morales, J. (2005) *Acta Palaeontol. Pol.* **50**, 219–238.
4. Gambaryan, P. P. (1974) *How Mammals Run: Anatomical Adaptations* (Wiley, New York).
5. Taylor, M. E. (1989) in *Carnivore Behavior, Ecology, and Evolution*, ed. Gittleman, J. L. (Cornell Univ. Press, Ithaca, New York), Vol. 1, pp. 382–409.
6. Davis, D. D. (1964) *Fieldiana Zool.* **3**, 1–339.
7. Chorn, J. & Hoffmann, R. S. (1978) *Mamm. Species* **110**, 1–6.
8. Roberts, M. S. & Gittleman, J. L. (1984) *Mamm. Species* **222**, 1–8.
9. Endo, H., Hayashi, Y., Yamagiwa, D., Kurohmaru, M., Koie, H., Yamaya, Y. & Kimura, J. (1999) *J. Anat.* **195**, 295–300.
10. Endo, H., Yamagiwa, D., Hayashi, Y., Koie, H., Yamaya, Y. & Kimura, J. (1999) *Nature* **397**, 309–310.
11. Ledje, C. & Arnason, U. (1996) *J. Mol. Evol.* **42**, 135–144.
12. Goldman, D., Rathna Giri, P. & O'Brien, S. J. (1989) *Evolution* **43**, 282–295.
13. Bininda-Emonds, O. R., Gittleman, J. L. & Purvis, A. (1999) *Biol. Rev. Camb. Philos. Soc.* **74**, 143–175.
14. Flynn, J. J., Finarelli, J. A., Zehr, S., Hsu, J. & Nedbal, M. A. (2005) *Syst. Biol.* **54**, 317–337.
15. Flynn, J. J., Nedbal, M. A., Dragoo, J. W. & Honeycutt, R. L. (2000) *Mol. Phylogenet. Evol.* **17**, 190–199.
16. Flynn, J. J. & Nedbal, M. A. (1998) *Mol. Phylogenet. Evol.* **9**, 414–426.
17. Ginsburg, L. (1999) in *The Miocene Land Mammals of Europe*, eds. Rössner, G.-E. & Heissig, K. (Pfeil, Munich), pp. 109–148.
18. O'Brien, S. J., Nash, W. G., Wildt, D. E., Bush, M. E. & Benveniste, R. E. (1985) *Nature* **317**, 140–144.
19. Wang, X. (1997) *J. Vert. Paleontol.* **17**, 184–198.
20. Wozencraft, W. C. (1989) in *Carnivore Behavior, Ecology, and Evolution*, ed. Gittleman, J.-L. (Cornell Univ. Press, Ithaca, New York), Vol. 1, pp. 495–535.
21. Wyss, A. R. & Flynn, J. J. (1993) in *Placentals, Mammal Phylogeny*, eds. Szalay, F., Novacek, M. & McKenna, M. (Springer, New York), pp. 32–52.
22. Vrana, P. B., Milinkovitch, M. C., Powell, J. R. & Wheeler, W. C. (1994) *Mol. Phylogenet. Evol.* **3**, 47–58.
23. Wayne, R. K., Benveniste, R. E., Janczewski, D. N. & O'Brien, S. J. (1989) in *Carnivore Behavior, Ecology, and Evolution*, ed. Gittleman, J. L. (Cornell Univ. Press, Ithaca, New York), Vol. 1, pp. 465–494.
24. Roussakis, S. J. (2001) *Senckenb. Lethaea* **81**, 347–358.
25. Baskin, J. (1998) in *Evolution of Tertiary Mammals of North America*, eds. Janis, C. M., Scott, K. M. & Jacobs, L. L. (Cambridge Univ. Press, Cambridge, U.K.), Vol. 1, pp. 144–151.
26. Ginsburg, L., Morales, J., Soria, D. & Herráez, E. (1997) *C. R. Acad. Sci.* **325**, 447–451.
27. Ginsburg, L., Maridet, O. & Mein, P. (2001) *Geodiversitas* **23**, 81–85.
28. Wallace, S. C. & Wang, X. (2004) *Nature* **431**, 556–559.
29. McClearn, D. (1992) *J. Mamm.* **73**, 245–261.
30. Ciro, E. (1992) Ph.D. thesis (Université de Poitiers, Poitiers, France).
31. Gould, S. J. (1980) *The Panda's Thumb* (Norton, New York).
32. de Beaumont, G. (1982) *Arch. Sci. Geneve* **35**, 143–152.